

Stem Cells All 'What Is...' Articles

Conditions Information

Abscess (Dental) Acid Reflux
Acne
Acoustic Neuroma (Vestibular Addison's Disease (Primary Ad
ADHD
Agoraphobia
Air Embolism (Gas Embolism)
Alcoholic Liver Disease (Alcoh
Alcoholism

Health Professional Sites

Ophthalmology

Urology

Other Navigation Links
About Us
News Licensing

Free Website Feeds
Free Tools & Content
Tell a Friend
Accessibility
Help / FAQ
Article Submission
Links
Contact Us

Clinical Trials



swallowed by children younger than 6 years came directly from a product, 29.8 percent were loose, and 8.2 percent were obtained from battery packaging. The most hazardous battery ingested, the 20-mm lithium cell, was intended for use in remote controls in 37.3 percent of cases. Study authors suggest that all consumer electronics powered by 20-mm lithium cells should require a secure battery compartment accessible with a tool (screwdriver) or child-resistant lock to prevent further pediatric ingestions. Parents must be vigilant, too, to prevent these ingestions.

Prc 01 Wh Syr 14 Fis In F 03

Source American Academy of Pediatrics

Please rate this article: Patient / Public: or **Health Professional:** (Hover over the stars then click to rate)



Any medical information published on this website is not intended as a substitute for informed medical advice and you should not take any action before consulting with a health care professional. For more information, please read our terms and conditions.

Contact Our News Editors

For any corrections of factual information, or to contact the editors please use our feedback form.

Please send any medical news or health news press releases to: pressrelease@medicalnewstoday.com

Back to top	Back to front pa	ge List of All Medical Articles
Privacy Policy	Terms and Conditions	© 2010 MediLexicon International Ltd



Lat

Fc

P

trar



Ads by Google

Battery Li Ion Severe Asthma BB Battery Asthma Allergy